

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

B28

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/335	A1	(11) International Publication Number: WO 97/44026 (43) International Publication Date: 27 November 1997 (27.11.97)
(21) International Application Number: PCT/US97/08866 (22) International Filing Date: 22 May 1997 (22.05.97) (30) Priority Data: 08/651,429 22 May 1996 (22.05.96) US (71) Applicant: NEUROMEDICA, INC. [US/US]; 99 Eric Street, Cambridge, MA 02139 (US). (72) Inventors: BRADLEY, Matthews, O.; 4309 Sundown Road, Laytonsville, MD 20882 (US). SHASHOUA, Victor, E.; 176 Tappan Street, Brookline, MA 02146 (US). WEBB, Nigel, L.; 1101 Green Valley Road, Bryn Mawr, PA 19010 (US). SWINDELL, Charles, S.; 613 Schiller Avenue, Merion, PA 19066 (US). (74) Agent: GATES, Edward, R.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: COMPOSITIONS COMPRISING CONJUGATES OF CIS-DOCOSAHEXAENOIC ACID AND TAXOTERE (57) Abstract This invention provides taxanes that are conjugates of <i>cis</i> -docosaheptaenoic acid and taxotere. The conjugates are useful in treating cancer.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMPOSITIONS COMPRISING CONJUGATES OF CIS-DOCOSAHEXAENOIC ACID AND TAXOTERE**Background of the Invention**

Taxol® (paclitaxel) was first isolated in 1971 from the bark of Taxus brevifolia and was
5 approved in 1992 by the US Food and Drug Administration for treatment of metastatic ovarian
cancer and later for breast cancer. Its mechanism of action is believed to involve promoting
formation and hyperstabilization of microtubules, thereby preventing the disassembly of
microtubules necessary for completion of cell division. It also has been reported that Taxol induces
expression of cytokines, affects the activity of kinases and blocks processes essential for metastasis,
10 in as yet uncharacterized mechanisms of action.


Taxol has attracted unusually strong scientific attention, not only because of its unique
antiproliferative mechanism of action, but also because it is active against nearly all cancers against
which it has been tested and because it has been discovered to be an analog of numerous closely
related compounds occurring naturally. These compounds, taxanes, are now recognized as a new
15 class of anticancer compounds.

Taxol's strength against cancers of diverse tissue origin also represents a significant
drawback. An ideal anticancer agent has tissue specificity, thereby reducing side-effects on normal
(dividing) cells. Taxol analogs with tissue specificity therefore are desired. Another drawback of
Taxol is its extreme insolubility. Taxol can be administered effectively in a solvent including
20 cremophor, which combination can provoke severe hypersensitive immune responses. As a result
of these drawbacks, and also as a result of the potential for modifying Taxol at numerous sites as
demonstrated by other naturally-occurring taxanes with anticancer activity, a search for more
selective taxanes was launched.

To date, more than 200 taxanes have been synthesized (or isolated) and tested *in vitro* or *in*
25 *vivo* for anticancer activity. The results, however, have been so disappointing that the National
Cancer Institute (NCI) generally no longer is interested in testing Taxol analogs. In general with
Taxol analogs, the solubility problems remain, and/or potency is sharply reduced, and/or selectivity
is not improved, and/or the ratio of the median toxic dose to the median effective dose ("therapeutic
index") is unacceptably reduced.


- 2 -

Taxol has the following formula:



Taxanes have the basic three ring structure (A, B and C), substituted or unsubstituted.

Taxol's carbons are numbered conventionally as follows:



Based upon the taxanes tested to date, as many questions have been raised as have been answered, and general rules have not been fashioned easily in predicting selectivity, activity and solubility. Firstly, no rules have emerged regarding selectivity. Those taxanes that are strongly active appear to have activity as broad as Taxol's activity, and no headway appears to have been made in terms of developing a more selective Taxol analog.

Some information about activity has emerged. Numerous substitutions have been made at C7, C9, C10, C19, R₁ and combinations thereof while retaining significant, but usually reduced, activity. Substitutions at C2, C4 and 2'OH, however, are generally not tolerated. These conclusions are only generalities, for example, because some substitutions at C9-C10 (cyclic derivatives) are not tolerated and some substitutions at C2 (meta substitutions on the phenyl) are tolerated. Likewise, the C13 side chain and, in particular, the 2'OH are required, although the minimum structural requirements of the side chain have not been calculated for therapeutic efficacy.

Attempts to improve Taxol's solubility have not resulted in successful clinical products. One approach has been to manufacture prodrugs of Taxol, which prodrugs undergo *in vivo* transformation into Taxol and some other product. Attempts were made to esterify the C7 hydroxy and 2' hydroxy groups, with the hope that the bond would be stable in solution (to permit preferred administration modes -i.v. over at least 24 hours) but would cleave readily *in vivo*. The groups tested were all hydrophilic and included amines, short carboxylic acids (using e.g. succinic anhydride and glutaric anhydride), sulfonic acids, amino acids and phosphates. Generally, activity was reduced although some success was obtained with certain derivatives. Again, no particular pattern emerged permitting one to predict reliably which groups could be substituted on Taxol to yield a therapeutically useful

- 3 -

product, although it was suggested that the 2' OH derivatives may cleave more easily than the C7 OH derivatives.


Several other factors add to the problem of predicting which Taxol analogs will be effective. Multiple mechanisms of action have been proposed in the literature, and a change in one position
5 may have no effect on activity on one such mechanism but may eliminate activity on another mechanism. In addition, changes that favorably influence activity may unfavorably influence bioavailability. For example, Taxol affects microtubule formation inside a cell, but a change in structure that increases intracellular activity may adversely affect the ability of Taxol to gain entry into a cell. Taxol also is known to bind to proteins, and the effect on activity that results from a
10 change in Taxol's binding to protein (in terms of conformation, cellular absorption and solubility) is unknown.

It has been reported that Taxol does not get into the brain, apparently excluded by the blood brain barrier. It is not known why this is so, as Taxol is lipophilic, gets into cells and might be expected to cross the blood brain barrier.

15 The most promising of the two hundred analogs tested is Taxotere (docetaxel), because of its slightly increased activity and solubility. Oddly, however, Taxotere differs from Taxol at sites which typically do not have a strong influence on activity, and one would not predict the improvements in Taxotere from these differences, even in hindsight.

Taxotere has the following formula:

20




25 DHA (docosahexaenoic acid) is a 22 carbon naturally-occurring, unbranched fatty acid that previously has been attached to drugs to help deliver them across the blood brain barrier. DHA is attached via the acid group to hydrophilic drugs and renders these drugs more hydrophobic (lipophilic). DHA is an important constituent of the brain and recently has been approved as an additive to infant formula. It is present in the milk of lactating women. The mechanism of action
30 by which DHA helps drugs conjugated to it cross the blood brain barrier is unknown.

Summary of the Invention


The present invention involves the unexpected finding that conjugates of Taxotere and a highly lipophilic group, a C22 unbranched carbon chain, have a different *selectivity* relative to Taxotere. The conjugates, in general, are believed to render the activity of the taxanes selective for colon cancer, breast cancer, and central nervous system cancer ("targeted cancers"). The conjugates, also unexpectedly, restrict the activity of the taxanes even within these three categories of cancer relative to that of Taxotere. The conjugates further unexpectedly, reduce sharply the activity of the taxanes relative to that of Taxotere in most cell lines of tissue types *other than* colon, breast, and central nervous system, thereby reducing potential side effects of the conjugates versus those of Taxotere. The therapeutic index of the conjugates is improved, versus that of Taxotere for targeted cancers.

According to one aspect of the invention, a composition of matter is provided. The composition is a covalent conjugate of *cis*-docosahexaenoic acid and Taxotere. Preferably, the conjugate consists of *cis*-docosahexaenoic acid and taxotere, wherein the *cis*-docosahexaenoic acid is conjugated directly to taxotere, free of a linker, for example via the carboxylic acid group of the *cis*-docosahexaenoic acid and a hydroxyl group of taxotere.

In one embodiment the conjugate is:




In another embodiment the conjugate is:




- 5 -

In another embodiment the conjugate is:



In another embodiment the conjugate is:



The conjugates of the invention can be isolated conjugates. An isolated conjugate is one which is separated from other different taxane conjugates.

Pharmaceutical preparations containing one or more of the foregoing conjugates also are provided. The pharmaceutical preparations preferably include a sterile, pharmaceutically acceptable carrier. The pharmaceutical preparations also can contain other anti-cancer agents.

The foregoing compositions of matter and pharmaceutical preparations are useful for treating cancer, preferably breast cancer, colon cancer and central nervous system cancer.

Brief Description of the Drawings

Figure 1 is a graph plotting concentration of conjugate 1 versus percent growth of leukemia cells.

Figure 2 is a graph plotting concentration of conjugate 1 versus percent growth of non-small cell lung cancer cells.

Figure 3 is a graph plotting concentration of conjugate 1 versus percent growth of colon cancer cells.

Figure 4 is a graph plotting concentration of conjugate 1 versus percent growth of CNS cancer cells.

Figure 5 is a graph plotting concentration of conjugate 1 versus percent growth of melanoma

cells.

Figure 6 is a graph plotting concentration of conjugate 1 versus percent growth of ovarian cancer cells.

Figure 7 is a graph plotting concentration of conjugate 1 versus percent growth of renal
5 cancer cells.

Figure 8 is a graph plotting concentration of conjugate 1 versus percent growth of prostate cancer cells.

Figure 9 is a graph plotting concentration of conjugate 1 versus percent growth of breast cancer cells.

10 Figure 10 is a graph plotting concentration of conjugate 2 versus percent growth of leukemia cells.

Figure 11 is a graph plotting concentration of conjugate 2 versus percent growth of non-small cell lung cancer cells.

Figure 12 is a graph plotting concentration of conjugate 2 versus percent growth of colon
15 cancer cells.

Figure 13 is a graph plotting concentration of conjugate 2 versus percent growth of CNS cancer cells.

Figure 14 is a graph plotting concentration of conjugate 2 versus percent growth of melanoma cells.

20 Figure 15 is a graph plotting concentration of conjugate 2 versus percent growth of ovarian cancer cells.

Figure 16 is a graph plotting concentration of conjugate 2 versus percent growth of renal cancer cells.

Figure 17 is a graph plotting concentration of conjugate 2 versus percent growth of prostate
25 cancer cells.

Figure 18 is a graph plotting concentration of conjugate 2 versus percent growth of breast cancer cells.

Figure 19 is a graph plotting concentration of Taxol versus percent growth of leukemia cells.

Figure 20 is a graph plotting concentration of Taxol versus percent growth of non-small cell
30 lung cancer cells.

Figure 21 is a graph plotting concentration of Taxol versus percent growth of colon cancer cells.

Figure 22 is a graph plotting concentration of Taxol versus percent growth of CNS cancer cells.

Figure 23 is a graph plotting concentration of Taxol versus percent growth of melanoma cells.

5 Figure 24 is a graph plotting concentration of Taxol versus percent growth of ovarian cancer cells.

Figure 25 is a graph plotting concentration of Taxol versus percent growth of renal cancer cells.

10 Figure 26 is a graph plotting concentration of Taxol versus percent growth of prostate cancer cells.

Figure 27 is a graph plotting concentration of Taxol versus percent growth of breast cancer cells.

Detailed Description of the Invention

cis-docosahexaenoic acid (DHA) is a naturally occurring fatty acid. It is an unbranched chain
15 fatty acid with six double bonds, all *cis*. Its structure is as follows:



DHA can be isolated, for example, from fish oil or can be chemically synthesized. These
20 methods, however, can generate *trans* isomers, which are difficult and expensive to separate and which may present safety problems in humans. The preferred method of production is biological synthesis to produce the all *cis* isomer. The preferred source of DHA is from Martek Biosciences Corporation of Columbia, Maryland. Martek has a patented system for manufacturing DHA using microalgae which synthesize only a single isomer of DHA, the all *cis* isomer. Martek's patents
25 include U.S. Pat. Nos. 5,374,657, 5,492,938, 5,407,957 and 5,397,591.

DHA also is present in the milk of lactating women, and Martek's licensee has obtained approval in Europe of DHA as a nutritional supplement for infant formula.

It is known that DHA can be unstable in the presence of oxygen. To stabilize DHA and its conjugates it is important to add anti-oxidants to the material after it is synthesized. One method of
30 stabilization is to make-up the newly synthesized material in the following solution: 100 g neat DHA-taxol plus 100 g of vehicle (100 ml propylene glycol, 70 mg alpha-tocopherol, 5 mg dilaurylthiodipropionic acid, 50 mg ascorbic acid) prepared and held under argon in amber, sealed


vials and stored at four degrees centigrade. The following anti-oxidants may also be employed: ascorbic acid, ascorbyl palmitate, dilauryl ascorbate, hydroquinone, butylated hydroxyanisole, sodium meta bisulfite, *l*- β carotene and α -tocopherol. A heavy metal chelator such as ethylenediamine tetra-acetic acid (EDTA) may also be used.

5 Paclitaxel was first isolated from the bark of Taxus brevifolia (Wani et al., J. Am. Chem. Soc., 93, 2325, 1971). Its isolation and synthesis have been reported extensively in the literature. Applicants obtained paclitaxel from a commercial source, Hauser Laboratories, of Boulder, Colorado.

Example 1

10 TAXOL

15



conjugate 1


To synthesize DHA-Taxol, a solution of Taxol (41 μ mol) in methylene chloride (2.5mL) under argon was mixed with 4-dimethylaminopyridine (41 μ mol), dicyclohexylcarbodiimide (82 μ mol), and DHA (41 μ mol) and the reaction mixture was stirred at ambient temperature for two hours. Following dilution with ether, the reaction mixture was washed with 5% hydrochloric acid, water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue produced 45mg (94%) of crystalline Taxol-DHA conjugate 1.

20


- 9 -

Example 2


TAXOL



A



B



conjugate 2

The production of conjugate 2 involves several steps including a number of protection-acylation-deprotection steps. A solution of Taxol (59 μ mol) in methylene chloride (2.5mL) was mixed at ambient temperature under argon with imidazole (147 μ mol) and triethylsilyl chloride (147 μ mol). The reaction mixture was stirred for thirty minutes, diluted with additional methylene chloride, washed with water, saturated aqueous sodium chloride, dried, and concentrated. Chromatography of the residue produced 50mg (88%) of intermediate A plus 5mg of the 2', 7-di(triethylsilyl) ether derivative. A solution of intermediate A (52 μ mol) in methylene chloride (3mL) was mixed at ambient temperature under argon with 4-dimethylaminopyridine (52 μ mol), dicyclohexylcarbodiimide (104 μ mol), and DHA (52 μ mol). The reaction mixture was stirred for ten hours, diluted with ether, passed through celite, and concentrated. Chromatography of the residue produced 65.9mg of intermediate B. A solution of intermediate B (51 μ mol) in acetonitrile (2mL) at 0°C under argon was mixed with 49% aqueous HF (0.2mL) and the reaction mixture was stirred for one hour. After dilution with ether, the reaction mixture was washed with water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue produced 44.6mg (75%) of Taxol-DHA conjugate 2.

- 10 -

Example 3

Conjugates 1 and 2 were sent to the United States National Cancer Institute (NCI) for screening in the NCI's anticancer screening program. The conjugates were provided in ethanol (approximately 40mg analog/2ml ethanol). The conjugates were sealed in vials under argon to avoid exposure of the conjugates to oxygen because the conjugates were believed to be sensitive to oxygen. Instructions were provided to store at 4°C and to open the vials only when ready for immediate experimental use. Instructions also were provided to use the ethanol solutions containing the conjugates directly or to dissolve the analogs further in DMSO (dimethylsulfoxide) at appropriate concentrations, with vortexing if necessary for adequate dispersal.

The activities of conjugates 1 and 2 were tested against 57 cancer cell lines. The results are presented in Figs. 1-9 for conjugate 1, Figs. 10-18 for conjugate 2 and Figs 19-27 for Taxol. To understand the data, reference is made to the guides provided by the NCI, excerpted as follows:

The Calculated Measurement of Effect: Percentage Growth (PG)

The measured effect of the compound on a cell line is currently calculated according to one or the other of the following two expressions:

If $(\text{Mean OD}_{\text{test}} - \text{Mean OD}_{\text{tzero}}) \geq 0$, then

$$\text{PG} = 100 \times (\text{Mean OD}_{\text{test}} - \text{Mean OD}_{\text{tzero}}) / (\text{Mean OD}_{\text{ctrl}} - \text{Mean OD}_{\text{tzero}})$$

If $(\text{Mean OD}_{\text{test}} - \text{Mean OD}_{\text{tzero}}) < 0$, then

$$\text{PG} = 100 \times (\text{Mean OD}_{\text{test}} - \text{Mean OD}_{\text{tzero}}) / \text{Mean OD}_{\text{tzero}}$$

Where:

Mean OD_{tzero} = The average of optical density measurements of SRB-derived color just before exposure of cells to the test compound.

Mean OD_{test} = The average of optical density measurements of SRB-derived color after 48 hours exposure of cells to the test compound.

Mean OD_{ctrl} = The average of optical density measurements of SRB-derived color after 48 hours with no exposure of cells to the test compound.

Experimental data was collected against each cell line. ... Each concentration is expressed as the \log_{10} (molar or $\mu\text{g/ml}$). ... The response parameters GI50, TGI, and LC50 are interpolated values representing the concentrations at which the PG is +50, 0, and -50, respectively. Sometimes these
5 response parameters cannot be obtained by interpolation. If, for instance, all of the PGs in a given row exceed +50, then none of the three parameters can be obtained by interpolation. In such a case, the value given for each response parameter is the highest concentration tested. ... This practice is extended similarly to the other possible situations where a response parameter cannot be obtained by interpolation.

10

Dose-Response Curves:

The dose-response curve page of the data package is created by plotting the PGs against the \log_{10} of the corresponding concentration for every cell line. The cell line curves are grouped by subpanel.
15 Horizontal lines are provided at the PG values of +50, 0, and -50. The concentrations corresponding to points where the curves cross these lines are the GI50, TGI and LC50, respectively.

Several important distinctions are apparent from the data. Most important, the patterns of anticancer activity for conjugates 1 and 2 differ from that of Taxol. In one sense, conjugates 1 and 2 are effective anticancer agents against a more restricted set of cancer cell lines. For example,
20 conjugates 1 and 2 were not very effective against any of the six leukemia cancer cell lines tested, whereas Taxol was somewhat effective against all four leukemia cell lines against which Taxol was tested. (See Figs. 1, 10 and 19.)

The relative activity against members within a class of cancers also was altered. For example, at TGI (horizontal line at zero in the graphs), Taxol was more effective against non-small
25 cell lung cancer line H522 than against H460 (by about 3 logs), whereas conjugates 1 and 2 were slightly more effective against H460 than H522. As another example, Taxol was least effective at TGI against CNSU251, whereas conjugate 1 was most effective against CNSU251 and conjugate 2 was also very effective against CNSU251 (relative to other CNS cell lines). As a further example, Taxol was equivalent in activity toward MDA-N and MDA-MB-435 breast cancer cell lines at all
30 concentrations tested, whereas conjugates 1 and 2 were more effective against MDA-N than MDA-MB-435 at all concentrations tested.

To further illustrate the differences in the activity of conjugates 1 and 2 versus that of Taxol,

the NCI subjected the data to a statistical analysis designed by the NCI to reflect differences in the pattern of activity of anticancer agents. Conjugate 1 and conjugate 2 were determined to be statistically different in their pattern of activity versus Taxol in this unique measurement by the NCI.

5 It also is to be noted that, in general, conjugates 1 and 2 were one thousand to ten thousand times less potent than Taxol for many cell lines tested. This reduction in activity is important, especially since conjugates 1 and 2 maintained strong activity against some cell lines. Conjugates 1 and 2 will be sufficiently active against certain cell lines, but will have, on average, a substantially and disproportionately lower activity against other cell lines, reducing potential side effects. For example, the TGI for Taxol against CNS SF-539 is -6.95, and the TGI for conjugate 1 against this cell line is -5.13 and for conjugate 2 is -5.53. (In other words, the activity of the analogs was reduced versus that of Taxol by less than 2 logs). The GI50 for Taxol against CNS SF 539 is -7.52, whereas the GI50s for conjugates 1 and 2 are -6.22 and -5.56, respectively (again less than 2 logs difference). In contrast, Taxol has a GI50 for cell line CNSSF 268 of less than -10.0, whereas conjugates 1 and 15 2 have GI50s for CNSSF 268 of 5.36 and 5.28, respectively. This represents a reduction of activity in the conjugates vs. that of Taxol by at least about 5 logs activity! On average, the GI50 for Taxol across all cell lines tested is at least -9.19. (It is probably much higher since concentrations less than -10 were not tested, and if Taxol was active at -10.0, -10 (instead of the actual lower value) was used in calculating the average of -9.19. There were 27 instances when this occurred.) The average 20 GI50s for conjugates 1 and 2, on the other hand, were 5.49 and 5.22, respectively. Therefore, the average difference in activity for Taxol vs. the conjugates is *at least* between 3 and 4 logs. Thus, the sharp reduction in the activity of the conjugates against many cell lines vs. a lesser reduction for other cell lines is expected to reduce the potential side effects of the conjugates versus those of Taxol at effective doses.

25 Cancers other than CNS, breast and colon cancer can be treated. For example, there was activity against non-small cell lung cancer cells, melanoma cells and ovarian cancer cells. However, the activity was relatively reduced and was extremely specific, limiting the utility of the conjugates for treating generally such cancers. In any event, cancer patients could be evaluated to determine if a conjugate is strongly active against the patient's cancer prior to selecting the conjugate as the 30 anti-cancer agent of choice for that patient.

The foregoing experiments establish that DHA conjugates have altered specificity versus that of Taxol for cancer cell lines. Because of this altered specificity, it also is clear that the conjugates


- 13 -

themselves are gaining access into the target cells (as opposed to simply releasing Taxol into the environment outside of the cell). Thus, the DHA moiety appears to selectively target certain cell types as opposed to others. The ability of the conjugates to gain entry into the targeted cells was unknown prior to the invention, and the ability of the DHA moiety to selectively target certain cell types was unexpected.

The same is true of DHA-Taxotere covalent conjugates, examples of which are presented below. Taxotere's synthesis has been reported extensively in the literature. One example is Kanazawa, A. et al., J. Organic Chem. 1994, Vol. 59, pp. 1238-1240.

Example 4


TAXOTERE



conjugate 3

A solution of Taxotere in methylene chloride under argon is mixed with 4-dimethylaminopyridine, dicyclohexylcarbodiimide, and DHA. The reaction mixture is stirred at ambient temperature. Radial chromatography of the residue is performed to produce Taxotere-DHA conjugate 3.


- 14 -

Example 5

- 25 A solution of Taxotere in dimethylformamide is mixed at ambient temperature under argon with imidazole and triethylsilyl chloride. The reaction mixture is stirred at ambient temperature, diluted with methylene chloride, washed with water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue is performed to produce intermediate C. A solution of intermediate C in methylene chloride is mixed at ambient temperature under argon with
- 30 4 dimethylaminopyridine, dicyclohexylcarbodiimide, and DHA. The reaction mixture is stirred at ambient temperature, diluted with ether, passed through celite, and concentrated. Radial chromatography of the residue is performed to produce intermediate D. A solution of intermediate

D in acetonitrile at 0°C under argon is mixed with 49% aqueous HF and the reaction mixture is stirred at the same temperature. After dilution with ether, the reaction mixture is washed with water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue is performed to produce Taxotere-DHA conjugate 4.

5


Example 6

A solution of Taxotere in dimethylformamide is mixed at ambient temperature under argon with imidazole and *tert*-butyldimethylsilyl chloride. The reaction mixture is stirred at ambient temperature, diluted with methylene chloride, washed with water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue is performed to produce intermediate E. A solution of intermediate E in methylene chloride is mixed at ambient temperature under argon with 4-dimethylaminopyridine, dicyclohexylcarbodiimide, and 1 equivalent of DHA. The reaction mixture is stirred at ambient temperature, diluted with ether, passed through celite, and concentrated. Radial chromatography of the residue is performed to produce intermediate F. (Intermediate H is also obtained and used in Example 8 below.) A solution of intermediate F in acetonitrile at 0°C under argon is mixed with aqueous HF and the reaction mixture is stirred at the same temperature. After dilution with ether, the reaction mixture is washed with water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue is performed

30

to produce Taxotere-DHA conjugate 4.


Example 7



- 17 -

intermediate G in acetonitrile at 0°C under argon is mixed with aqueous HF and the reaction mixture is stirred at the same temperature. After dilution with ether, the reaction mixture is washed with water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue is performed to produce Taxotere-DHA conjugate 5.

5

Example 8

25

A solution of taxotere in dimethylformamide is mixed at ambient temperature under argon with imidazole and *tert*-butyldimethylsilyl chloride. The reaction mixture is stirred at ambient temperature, diluted with methylene chloride, washed with water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue is performed to produce intermediate E. A solution of intermediate E in methylene chloride is mixed at ambient temperature under argon with 4-dimethylaminopyridine, dicyclohexylcarbodiimide, and 1 equivalent of DHA. The reaction mixture is stirred at ambient temperature, diluted with ether, passed through celite, and concentrated. Radial chromatography of the residue is performed to produce intermediate H and

30

intermediate F which was used above in Example 6. A solution of intermediate H in acetonitrile at 0°C under argon is mixed with aqueous HF and the reaction mixture is stirred at the same temperature. After dilution with ether, the reaction mixture is washed with water, saturated aqueous sodium chloride, dried, and concentrated. Radial chromatography of the residue is performed to produce Taxotere-DHA conjugate 6.

The compounds useful in the invention may be delivered in the form of anti-cancer cocktails. An anti-cancer cocktail is a mixture of any one of the compounds useful with this invention with another anti-cancer agent such as an anti-cancer drug, a cytokine, and/or supplementary potentiating agent(s). The use of cocktails in the treatment of cancer is routine. In this embodiment, a common administration vehicle (e.g., pill, tablet, implant, injectable solution, etc.) would contain both the conjugate useful in this invention and the anti-cancer drug and/or supplementary potentiating agent.

Anti-cancer agents include anti-cancer drugs. Anti-cancer drugs are well known and include: Acivicin; Aclarubicin; Acodazole Hydrochloride; Acronine; Adozelesin; Aldesleukin; Altretamine; Ambomycin; Ametantrone Acetate; Aminoglutethimide; Amsacrine; Anastrozole; Anthramycin; Asparaginase; Asperlin; Azacitidine; Azetepa; Azotomycin; Batimastat; Benzodepa; Bicalutamide; Bisantrene Hydrochloride; Bisnafide Dimesylate; Bizelesin; Bleomycin Sulfate; Brequinar Sodium; Bropirimine; Busulfan; Cactinomycin; Calusterone; Caracemide; Carbetimer; Carboplatin; Carmustine; Carubicin Hydrochloride; Carzelesin; Cedefingol; Chlorambucil; Cirolemycin; Cisplatin; Cladribine; Crisnatol Mesylate; Cyclophosphamide; Cytarabine; Dacarbazine; Dactinomycin; Daunorubicin Hydrochloride; Decitabine; Dexormaplatin; Dezaguanine; Dezaguanine Mesylate; Diaziquone; Docetaxel; Doxorubicin; Doxorubicin Hydrochloride; Droloxifene; Droloxifene Citrate; Dromostanolone Propionate; Duazomycin; Edatrexate; Eflornithine Hydrochloride; Elsamitrucin; Enloplatin; Enpromate; Epipropidine; Epirubicin Hydrochloride; Erbulozole; Erorubicin Hydrochloride; Estramustine; Estramustine Phosphate Sodium; Etanidazole; Etoposide; Etoposide Phosphate; Etoprine; Fadrozole Hydrochloride; Fazarabine; Fenretinide; Floxuridine; Fludarabine Phosphate; Fluorouracil; Flurocitabine; Fosquidone; Fostriccin Sodium; Gemcitabine; Gemcitabine Hydrochloride; Hydroxyurea; Idarubicin Hydrochloride; Ifosfamide; Ilmofofosine; Interferon Alfa-2a; Interferon Alfa-2b; Interferon Alfa-n1; Interferon Alfa-n3; Interferon Beta-1 a; Interferon Gamma-1 b; Iproplatin; Irinotecan Hydrochloride; Lanreotide Acetate; Letrozole; Leuprolide Acetate; Liarozole Hydrochloride; Lometrexol Sodium; Lomustine; Losoxantrone Hydrochloride; Masoprocil; Maytansine; Mechlorethamine Hydrochloride; Megestrol Acetate; Melengestrol Acetate; Melphalan; Menogaril; Mercaptopurine;

Methotrexate; Methotrexate Sodium; Metoprine; Meturedopa; Mitindomide; Mitocarcin; Mitocromin; Mitogillin; Mitomalcin; Mitomycin; Mitosper; Mitotane; Mitoxantrone Hydrochloride; Mycophenolic Acid; Nocodazole; Nogalamycin; Ormaplatin; Oxisuran; Paclitaxel; Pegaspargase; Peliomycin; Pentamustine; Peplomycin Sulfate; Perfosfamide; Pipobroman; Pipsulfan;
 5 Piroxantrone Hydrochloride; Plicamycin; Plomestane; Porfimer Sodium; Porfiromycin; Prednimustine; Procarbazine Hydrochloride; Puromycin; Puromycin Hydrochloride; Pyrazofurin; Riboprime; Rogletimide; Safingol; Safingol Hydrochloride; Semustine; Simtrazene; Sparfosate Sodium; Sparsomycin; Spirogermanium Hydrochloride; Spiromustine; Spiroplatin; Streptonigrin; Streptozocin; Sulofenur; Talisomycin; Tecogalan Sodium; Tegafur; Teloxantrone Hydrochloride;
 10 Temoporfin; Teniposide; Teroxirone; Testolactone; Thiamiprine; Thioguanine; Thiotepa; Tiazofurin; Tirapazamine; Topotecan Hydrochloride; Toremifene Citrate; Trestolone Acetate; Tricirbine Phosphate; Trimetrexate; Trimetrexate Glucuronate; Triptorelin; Tubulozole Hydrochloride; Uracil Mustard; Uredepa; Vapreotide; Verteporfin; Vinblastine Sulfate; Vincristine Sulfate; Vindesine; Vindesine Sulfate; Vinepidine Sulfate; Vinglycinat Sulfate; Vinleurosine Sulfate; Vinorelbine
 15 Tartrate; Vinrosidine Sulfate; Vinzolidine Sulfate; Vorozole; Zaniplatin; Zinostatin; Zorubicin Hydrochloride.

Other anti-cancer drugs include: 20-epi-1,25 dihydroxyvitamin D₃; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine;
 20 anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; anti-dorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III
 25 derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; canarypox IL-2; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700;
 30 cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cctrorcelix; chlorins; chloroquinoxaline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin

analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatan; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydrodidemnin B; deslorelin; dexifosfamide; dextrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine;

5 dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docosanol; dolasetron; doxifluridine; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine;

10 fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofofosine; ilomastat; imidazoacridones; imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists;

15 interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; irinotecan; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide + estrogen + progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum

20 compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; lovastatin; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocil; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mismatched double stranded RNA; mitoguazone; mitolactol;

25 mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotenc; molgramostim; monoclonal antibody, human chorionic gonadotrophin; monophosphoryl lipid A + myobacterium cell wall sk; mopidamol; multiple drug resistance gene inhibitor; multiple tumor suppressor 1-based therapy; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin;

30 nagrestip; naloxone + pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; neutral endopeptidase; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; O6-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone;

ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; 5 phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; 10 purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rogletimide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; 15 senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; signal transduction modulators; single chain antigen binding protein; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stem cell inhibitor; stem-cell division inhibitors; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; synthetic glycosaminoglycans; tallimustine; 20 tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; temozolomide; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thalidomide; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrigan; thyroid stimulating hormone; tin ethyl etiopurpurin; 25 tirapazamine; titanocene dichloride; topotecan; topsentin; toremifene; totipotent stem cell factor; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; vector system, erythrocyte gene therapy; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; 30 vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; zinostatin stimalamer.

Supplementary potentiating agents likewise are well characterized and include: Tricyclic anti-depressant drugs (e.g., imipramine, desipramine, amitriptyline, clomipramine, trimipramine,

doxepin, nortriptyline, protriptyline, amoxapine and maprotiline); non-tricyclic anti-depressant drugs (e.g., sertraline, trazodone and citalopram); Ca⁺⁺ antagonists (e.g., verapamil, nifedipine, nitrendipine and caroverine); Calmodulin inhibitors (e.g., prenylamine, trifluoroperazine and clomipramine); Amphotericin B; Triparanol analogues (e.g., tamoxifen); antiarrhythmic drugs (e.g., quinidine);
5 antihypertensive drugs (e.g., reserpine); Thiol depleters (e.g., buthionine and sulfoximine) and Multiple Drug Resistance reducing agents such as Cremaphor EL.

The compounds of the invention also can be administered with cytokines such as granulocyte colony stimulating factor.

The compounds of the invention, when used in cocktails, are administered in therapeutically
10 effective amounts. A therapeutically effective amount will be determined by the parameters discussed below; but, in any event, is that amount which establishes a level of the drug(s) in the area of the tumor which is effective in inhibiting the tumor growth.

When administered, the formulations of the invention are applied in pharmaceutically acceptable amounts and in pharmaceutically acceptable compositions. Such preparations may
15 routinely contain salts, buffering agents, preservatives, compatible carriers, and optionally other therapeutic ingredients. When used in medicine the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically acceptable salts thereof and are not excluded from the scope of the invention. Such pharmacologically and pharmaceutically acceptable salts include, but are not limited to, those
20 prepared from the following acids: hydrochloric, hydrobromic, sulphuric, nitric, phosphoric, maleic, acetic, salicylic, p-toluene sulfonic, tartaric, citric, methane sulfonic, formic, malonic, succinic, naphthalene-2-sulfonic, and benzene sulfonic. Also, pharmaceutically acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts.

Suitable buffering agents include: acetic acid and a salt (1-2% W/V); citric acid and a salt
25 (1-3% W/V); boric acid and a salt (0.5-2.5% W/V); and phosphoric acid and a salt (0.8-2% W/V).

Suitable preservatives include benzalkonium chloride (0.003-0.03% W/V); chlorobutanol (0.3-0.9% W/V); parabens (0.01-0.25% W/V) and thimerosal (0.004-0.02% W/V).

The active compounds of the present invention may be a pharmaceutical composition having a therapeutically effective amount of a conjugate of the invention optionally included in a
30 pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier" as used herein means one or more compatible solid or liquid filler, dilutants or encapsulating substances which are suitable for administration to a human or other animal. The term "carrier" denotes an organic or

inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions are capable of being commingled with the molecules of the present invention, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficacy.

5 Compositions suitable for parenteral administration conveniently comprise a sterile preparation of the conjugates of the invention. This preparation may be formulated according to known methods. Formulations for taxanes can be found in Chapter 9 of Taxol: Science and Applications, CRC Press, Inc., 2000 Corporate Boulevard, N.W., Boca Raton, FL 33431. In general, Taxol has been formulated as a 6 mg/ml cremophor EL (polyoxyethylated castor oil)/ethanol
10 mixture, which is diluted to final volume with normal saline or 5% dextrose. A 15mg/ml solution of taxotere has been formulated in polysorbate 80 (polyoxyethylene sorbitanmonooleate)/ethanol mixture, diluted with 5% dextrose.

The sterile preparation thus may be a sterile solution or suspension in a non-toxic parenterally-acceptable diluent or solvent. In addition, sterile, fixed oils are conventionally
15 employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono ordi-glycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables. Carrier formulations suitable for oral, subcutaneous, intravenous, intramuscular, etc. can be found in Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, PA.

20 The invention is used in connection with treating subjects having, suspected of having, developing or suspected of developing cancer. A subject as used herein means humans, primates, horses, cows, pigs, sheep, goats, dogs, cats and rodents.

The conjugates of the invention are administered in effective amounts. An effective amount means that amount necessary to delay the onset of, inhibit the progression of or halt altogether the
25 onset or progression of the particular condition being treated. In general, an effective amount will be that amount necessary to inhibit mammalian cancer cell proliferation in-situ. When administered to a subject, effective amounts will depend, of course, on the particular condition being treated; the severity of the condition; individual patient parameters including age, physical condition, size and weight; concurrent treatment; frequency of treatment; and the mode of administration. These factors
30 are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is preferred generally that a maximum dose be used, that is, the highest safe dose according to sound medical judgment.

Dosage may be adjusted appropriately to achieve desired drug levels, locally or systemically. Generally, daily oral doses of active compounds will be from about 0.01 mg/kg per day to 1000 mg/kg per day. It is expected that IV doses in the range of about 1 to 1000 mg/m² per day will be effective. In the event that the response in a subject is insufficient at such doses, even higher doses
5 (or effective higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits. Continuous IV dosing over, for example 24 hours or multiple doses per day are contemplated to achieve appropriate systemic levels of compounds.

A variety of administration routes are available. The particular mode selected will depend of course, upon the particular drug selected, the severity of the disease state being treated and the
10 dosage required for therapeutic efficacy. The methods of this invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of the active compounds without causing clinically unacceptable adverse effects. Such modes of administration include oral, rectal, sublingual, topical, nasal, transdermal or parenteral routes. The term "parenteral" includes subcutaneous, intravenous, intramuscular, or
15 infusion. Intravenous routes are preferred.

The compositions may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the conjugates of the invention into association with a carrier which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately
20 bringing the compounds into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product.

Compositions suitable for oral administration may be presented as discrete units such as capsules, cachets, tablets, or lozenges, each containing a predetermined amount of the active compound. Other compositions include suspensions in aqueous liquors or non-aqueous liquids such
25 as a syrup, an elixir, or an emulsion.

Other delivery systems can include time-release, delayed release or sustained release delivery systems. Such systems can avoid repeated administrations of the active compounds of the invention, increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer based systems such
30 as polylactic and polyglycolic acid, polyanhydrides and polycaprolactone; nonpolymer systems that are lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono-, di and triglycerides; hydrogel release systems; silastic systems; peptide based systems;

wax coatings, compressed tablets using conventional binders and excipients, partially fused implants and the like. In addition, a pump-based hardware delivery system can be used, some of which are adapted for implantation.

5 A long-term sustained release implant also may be used. "Long-term" release, as used herein, means that the implant is constructed and arranged to deliver therapeutic levels of the active ingredient for at least 30 days, and preferably 60 days. Long-term sustained release implants are well known to those of ordinary skill in the art and include some of the release systems described above. Such implants can be particularly useful in treating solid tumors by placing the implant near or directly within the tumor, thereby affecting localized, high-doses of the compounds of the
10 invention.


The analogs of the invention also are useful, in general, for treating mammalian cell proliferative disorders other than cancer, including psoriasis, actinic keratosis, etc.

Those skilled in the art will be able to recognize with no more than routine experimentation numerous equivalents to the specific products and processes described above. Such equivalents are
15 intended to be included within the scope of the appended claims.

We claim as follows:


CLAIMS

1. A composition of matter comprising
a covalent conjugate of *cis*-docosahexaenoic acid and Taxotere.
- 5 2. The composition of claim 1, wherein the *cis*-docosahexaenoic acid is conjugated directly to Taxotere.
3. The composition of matter of claim 1, wherein the composition is:




15

4. The composition of matter of claim 1, wherein the composition is:




25

5. The composition of matter of claim 1, wherein the composition is:



6. The composition of matter of claim 1, wherein the composition is:




7. A pharmaceutical composition comprising
a conjugate of *cis*-docosahexaenoic acid and Taxotere, and
10 a sterile, pharmaceutically acceptable carrier.

8. The pharmaceutical composition of claim 7, wherein the *cis*-docosahexaenoic acid is
conjugated directly to Taxotere.


15

9. The pharmaceutical composition of claim 7, wherein the:




25


10. The pharmaceutical composition of claim 7, wherein the conjugate is:



11. The pharmaceutical composition of claim 7, wherein the conjugate is:





12. The pharmaceutical composition of claim 7, wherein the conjugate is:







- 15 13. The pharmaceutical composition of claim 7 further comprising an anti-cancer agent other than the conjugate.
14. The pharmaceutical of claim 13 wherein the anti-cancer agent is selected from the group consisting of:



20 Aminoglutethimide; Asparaginase; Bleomycin; L-Buthiamine Sulfoxide; Busulfan; Camptothecin; Carboplatin; Carmustine (BCNU); Chlorambucil; Cisplatin (cis-DDP); Cyclophosphamide; Cytarabine HCl; Dacarbazine; Dactinomycin; Daunorubicin HCl; Doxorubicin HCl; Edatrexate; Estramustine phosphate sodium; Etoposide (V16-213); Floxuridine; Fluorouracil (5-FU); Flutamide; Gallium Nitrite; Hydroxyurea (hydroxycarbamide); Idarubicin; Ifosfamide; Interferon Alfa-2a, Alfa 2b; Leuprolide acetate (LHRH-releasing factor analogue); Lomustine (CCNU); Mechlorethamine HCl (nitrogen mustard); Megestrol; melphalan; Mercaptopurine; Methotrexate (MTX); Mitomycin; Mitotane (o,p'-DDD); Mitoxantrone HCl; Octreotide; Plicamycin; Prednisone; Procarbazine HCl; Streptozocin; Tamoxifen citrate; Taxanes; Taxoids; Thioguanine; Thiotepa; Tiasofuran; Topotecan; Vinblastine sulfate; Vincristine sulfate; Amsacrine (m-AMSA); Azacitidine;



25 Hexamethylmelamine (HMM); Interleukin 2; Mitoguazone (methyl-GAG); Methyl glyoxal bis-guanyldiazide (MGBG); Pentostatin; Semustine (methyl-CCNU); and Teniposide (VM-26).



**Fig. 1****Fig. 2**



**Fig. 3****Fig. 4**

**Fig. 5****Fig. 6**

**Fig. 7****Fig. 8**

**Fig. 9****Fig. 10**

**Fig. 11****Fig. 12**

**Fig. 13****Fig. 14**

10/10/10



10/10/10

10/10/10



10/10/10



10/10/10



10/10/10



**Fig. 15****Fig. 16**



9/14

**Fig. 17****Fig. 18**


**Fig. 19****Fig. 20**

**Fig. 21****Fig. 22**

**Fig. 23****Fig. 24**

**Fig. 25****Fig. 26**

14/14

*Fig. 27*

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/08866

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/335

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	No relevant documents disclosed -----	

☐

Further documents are listed in the continuation of box C.

☐

Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

13 August 1997

Date of mailing of the international search report

03.09.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Economou, D

